

Answers to Exercises with Matrices

1.
$$\begin{bmatrix} 11 & 22 \\ 33 & 44 \\ 55 & 66 \end{bmatrix}$$

2. $1 \cdot 3 + 2 \cdot 4 = 11$

3. $[11 \ 2]$

4.
$$\begin{bmatrix} 11 & 2 \\ 110 & 20 \end{bmatrix}$$

5. no answer – can't multiply a 1x3 matrix by a 2x1 matrix

6.
$$\begin{bmatrix} 3 & 6 & 21 \\ 4 & 8 & 28 \end{bmatrix}$$

7.
$$\begin{bmatrix} 18 & 21 & 24 \\ 5 & 10 & 15 \end{bmatrix}$$

8.
$$\begin{bmatrix} 7 & 8 & 9 \\ 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix}$$

9.
$$\begin{bmatrix} 5 & 10 \\ 15 & 20 \\ 25 & 30 \end{bmatrix}$$

10.
$$\begin{bmatrix} -1 & -2 \\ -3 & -4 \\ 5 & 6 \end{bmatrix}$$

11.
$$\begin{bmatrix} -9 & -18 \\ -27 & -36 \\ -45 & -54 \end{bmatrix}$$

12.
$$\begin{bmatrix} 1 & 3 & 5 \\ 2 & 4 & 6 \end{bmatrix}$$

13.
$$\begin{bmatrix} 1 & 3 & 5 \\ 2 & 4 & 6 \end{bmatrix}^T$$
 (the superscript T means transpose, same as in question 12)

14.
$$\begin{bmatrix} 1 & 2 \\ 3 & 4 \\ 5 & 6 \end{bmatrix}$$

15. $x=0$ and $y=1$

16. $x=5$ and $y=2$

17.
$$\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$
, since the identity matrix times itself equals the identity matrix.

18. Let $\begin{bmatrix} a & b \\ c & d \end{bmatrix} = \begin{bmatrix} 1 & 2 \\ -3 & 4 \end{bmatrix}^{-1}$, so $\begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} 1 & 2 \\ -3 & 4 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$ by definition of an inverse.

Hence $\begin{bmatrix} 1a-3b & 2a+4b \\ c-3d & 2c+4d \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$. Solve $1a-3b=1$ and $2a+4b=0$ to get

$a = \frac{2}{5}$, $b = \frac{-1}{5}$. Solve $c-3d=0$ and $2c+4d=1$ to get $c = \frac{3}{10}$, $d = \frac{1}{10}$. Hence the answer is

$\begin{bmatrix} 0.4 & -0.2 \\ 0.3 & 0.1 \end{bmatrix}$. Verify this! $\begin{bmatrix} 1 & 2 \\ -3 & 4 \end{bmatrix} \begin{bmatrix} 0.4 & -0.2 \\ 0.3 & 0.1 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$, so indeed

$$\begin{bmatrix} 1 & 2 \\ -3 & 4 \end{bmatrix}^{-1} = \begin{bmatrix} 0.4 & -0.2 \\ 0.3 & 0.1 \end{bmatrix}.$$

19. You can check the rules for determinants in the text:

- $1 \cdot 4 - 3 \cdot 2 = -2$, not singular
- 0, singular
- -30 , not singular
- -30 , not singular

20. square matrix

21. column matrix or column vector (or just “vector”)

22. row matrix or row vector

23. identity matrix

24. diagonal matrix

25. symmetric matrix

26. permutation matrix

27. upper triangular matrix

28. lower triangular matrix

$$29. \begin{bmatrix} \frac{1}{n} & \frac{1}{n} & \dots & \frac{1}{n} \\ \frac{1}{n} & \frac{1}{n} & \dots & \frac{1}{n} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{1}{n} & \frac{1}{n} & \dots & \frac{1}{n} \end{bmatrix}$$

$$30. \begin{bmatrix} \frac{1}{n} \sum x_i \\ \frac{1}{n} \sum x_i \\ \vdots \\ \frac{1}{n} \sum x_i \end{bmatrix} = \begin{bmatrix} \bar{x} \\ \bar{x} \\ \vdots \\ \bar{x} \end{bmatrix} \text{ where } \bar{x} \text{ is the mean of } x_1 \text{ through } x_n$$

$$31. \begin{bmatrix} x_1 - \bar{x} \\ x_2 - \bar{x} \\ \vdots \\ x_n - \bar{x} \end{bmatrix}$$

$$32. \left[\mathbf{I} - \frac{1}{n} \mathbf{ii}' \right] \mathbf{x} = \mathbf{Ix} - \frac{1}{n} \mathbf{ii}' \mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} - \begin{bmatrix} \bar{x} \\ \bar{x} \\ \vdots \\ \bar{x} \end{bmatrix} = \begin{bmatrix} x_1 - \bar{x} \\ x_2 - \bar{x} \\ \vdots \\ x_n - \bar{x} \end{bmatrix}$$

$$33. \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 0 \end{bmatrix}$$

$$34. \mathbf{M} \equiv \left[\mathbf{I} - \frac{1}{n} \mathbf{ii}' \right] = \begin{bmatrix} 1 - \frac{1}{n} & -\frac{1}{n} & \cdots & -\frac{1}{n} \\ -\frac{1}{n} & 1 - \frac{1}{n} & \cdots & -\frac{1}{n} \\ \vdots & \vdots & \ddots & \vdots \\ -\frac{1}{n} & -\frac{1}{n} & \cdots & 1 - \frac{1}{n} \end{bmatrix}$$

$$35. \sum x_i$$

$$36. \mathbf{i}'(\mathbf{Mx}) = \sum_{i=1}^n x_i - n\bar{x} = n\bar{x} - n\bar{x} = 0 \quad (\text{for } \mathbf{Mx}, \text{ plug in your four-previous answer})$$

$$37. \mathbf{M}$$

$$38. \sum x_i^2$$

$$39. \sum (x_i - \bar{x})^2$$

$$40. \sum x_i y_i$$

$$41. \sum (x_i - \bar{x})(y_i - \bar{y})$$

$$42. \frac{1}{n-1} \sum (x_i - \bar{x})(y_i - \bar{y})$$

$$43. \begin{bmatrix} x_{11} - \bar{x}_1 & x_{12} - \bar{x}_2 & \cdots & x_{1k} - \bar{x}_k \\ x_{21} - \bar{x}_1 & x_{22} - \bar{x}_2 & \cdots & x_{2k} - \bar{x}_k \\ \vdots & \vdots & \ddots & \vdots \\ x_{n1} - \bar{x}_1 & x_{n2} - \bar{x}_2 & \cdots & x_{nk} - \bar{x}_k \end{bmatrix} \quad \text{where } \bar{x}_i \text{ is the mean of } x_{i1} \text{ through } x_{ni}$$

$$44. \begin{bmatrix} \sum (x_{i1} - \bar{x}_1)^2 & \sum (x_{i1} - \bar{x}_1)(x_{i2} - \bar{x}_2) & \cdots & \sum (x_{i1} - \bar{x}_1)(x_{ik} - \bar{x}_k) \\ \sum (x_{i2} - \bar{x}_2)(x_{i1} - \bar{x}_1) & \sum (x_{i2} - \bar{x}_2)^2 & \cdots & \sum (x_{i2} - \bar{x}_2)(x_{ik} - \bar{x}_k) \\ \vdots & \vdots & \ddots & \vdots \\ \sum (x_{ik} - \bar{x}_k)(x_{i1} - \bar{x}_1) & \sum (x_{ik} - \bar{x}_k)(x_{i2} - \bar{x}_2) & \cdots & \sum (x_{ik} - \bar{x}_k)^2 \end{bmatrix}$$

$$45. \begin{bmatrix} \text{var}(x_{.1}) & \text{cov}(x_{.1}, x_{.2}) & \cdots & \text{cov}(x_{.1}, x_{.k}) \\ \text{cov}(x_{.2}, x_{.1}) & \text{var}(x_{.2}) & \cdots & \text{cov}(x_{.2}, x_{.k}) \\ \vdots & \vdots & \ddots & \vdots \\ \text{cov}(x_{.k}, x_{.1}) & \text{cov}(x_{.k}, x_{.2}) & \cdots & \text{var}(x_{.k}) \end{bmatrix} \quad \text{where}$$

$$\text{cov}(x_{.a}, x_{.b}) = \frac{1}{n-1} \sum (x_{ia} - \bar{x}_a)(x_{ib} - \bar{x}_b) \quad \text{and} \quad \text{var}(x_{.a}) = \text{cov}(x_{.a}, x_{.a})$$

$$46. \mathbf{I}$$

47. **I**48. **C**49. **I**50. **D'C'**51. **D⁻¹C⁻¹**52. **A⁻¹A' = A⁻¹A = I**, since **A** is symmetric53. **C⁻¹C'** (this cannot be simplified, since **C** is not necessarily symmetric)54. **B⁻¹(AB)'A⁻¹ = B⁻¹B'A'A⁻¹ = B⁻¹BAA⁻¹ = II = I**

55.
$$\begin{bmatrix} x_1 & x_2 \end{bmatrix} \begin{bmatrix} 1 & 1.5 \\ 1.5 & 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

56.
$$\begin{bmatrix} x_1 & x_2 & x_3 \end{bmatrix} \begin{bmatrix} 0 & 0.5 & 0.5 \\ 0.5 & 0 & 0.5 \\ 0.5 & 0.5 & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}$$

57. not a quadratic form – the term $7x_1$ is not quadratic

58.
$$\begin{bmatrix} x_1 \end{bmatrix} \begin{bmatrix} 1 \end{bmatrix} \begin{bmatrix} x_1 \end{bmatrix}$$

59.
$$\begin{bmatrix} x_1 & x_2 \end{bmatrix} \begin{bmatrix} 0 & 0.5 \\ 0.5 & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

60. not a quadratic form

61. not a quadratic form

62.
$$\begin{bmatrix} x_1 & x_2 & x_3 \end{bmatrix} \begin{bmatrix} 3 & 0.5 & -5/2\pi \\ 0.5 & 1 & -6 \\ -5/2\pi & -6 & 7 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}$$

63.
$$\begin{bmatrix} x_1 & x_2 & x_3 \end{bmatrix} \begin{bmatrix} \alpha & \delta/2\rho & \phi/2 \\ \delta/2\rho & \beta & \sigma/2 \\ \phi/2 & \sigma/2 & \gamma \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}$$

64. The answer to the first step is $a_1u_1 + a_2u_2 + \dots + a_nu_n$. The answer to the second step is $\frac{\partial \mathbf{a}'\mathbf{u}}{\partial u_i} = \frac{\partial (a_1u_1 + a_2u_2 + \dots + a_nu_n)}{\partial u_i} = a_i$. The answer to the third step is therefore

$$\begin{bmatrix} a_1 \\ \vdots \\ a_n \end{bmatrix} = \begin{bmatrix} a_1 & \dots & a_n \end{bmatrix}'. \text{ By definition, } \frac{\partial f(\mathbf{u})}{\partial \mathbf{u}} = \begin{bmatrix} \frac{\partial f(\mathbf{u})}{\partial u_1} & \dots & \frac{\partial f(\mathbf{u})}{\partial u_n} \end{bmatrix}', \text{ so applying}$$

this formula to the function $f(\mathbf{u}) = \mathbf{a}'\mathbf{u}$, you see that you have proved the formula

$$\frac{\partial (\mathbf{a}'\mathbf{u})}{\partial \mathbf{u}} = \mathbf{a}.$$

65. First compute the scalar $\mathbf{u}'\mathbf{A}\mathbf{u}$, and compute its vector of derivatives. Then compute $2\mathbf{A}\mathbf{u}$ and see if you get the same answer.

$$\begin{aligned} \mathbf{u}'\mathbf{A}\mathbf{u} &= \begin{bmatrix} u_1 & u_2 \end{bmatrix} \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix} \begin{bmatrix} u_1 \\ u_2 \end{bmatrix} \\ &= \begin{bmatrix} a_{11}u_1 + a_{21}u_2 & a_{12}u_1 + a_{22}u_2 \end{bmatrix} \begin{bmatrix} u_1 \\ u_2 \end{bmatrix} \\ &= a_{11}u_1^2 + a_{21}u_1u_2 + a_{12}u_1u_2 + a_{22}u_2^2 \end{aligned}$$

$$\frac{\partial(\mathbf{u}'\mathbf{A}\mathbf{u})}{\partial\mathbf{u}} = \begin{bmatrix} \frac{\partial(\mathbf{u}'\mathbf{A}\mathbf{u})}{\partial u_1} \\ \frac{\partial(\mathbf{u}'\mathbf{A}\mathbf{u})}{\partial u_2} \end{bmatrix} = \begin{bmatrix} 2a_{11}u_1 + (a_{21} + a_{12})u_2 \\ (a_{21} + a_{12})u_1 + 2a_{22}u_2 \end{bmatrix} \quad (1)$$

$$2\mathbf{A}\mathbf{u} = 2 \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix} \begin{bmatrix} u_1 \\ u_2 \end{bmatrix} = \begin{bmatrix} 2a_{11}u_1 + 2a_{12}u_2 \\ 2a_{21}u_1 + 2a_{22}u_2 \end{bmatrix} \quad (2)$$

Comparing (1) and (2), it is apparent that they are equal if $a_{21} + a_{12} = 2a_{12}$ and $a_{21} + a_{12} = 2a_{21}$. This can be true only if $a_{12} = a_{21}$. Thus (1) and (2) are equal if and

only if \mathbf{A} is symmetric. Thus $\frac{\partial(\mathbf{u}'\mathbf{A}\mathbf{u})}{\partial\mathbf{u}} = 2\mathbf{A}\mathbf{u}$ if and only if \mathbf{A} is symmetric.

$$66. \frac{\partial \mathbf{f}}{\partial \mathbf{u}'} = \begin{bmatrix} \frac{\partial f_1(\mathbf{u})}{\partial u_1} & \frac{\partial f_1(\mathbf{u})}{\partial u_2} & \frac{\partial f_1(\mathbf{u})}{\partial u_3} \\ \frac{\partial f_2(\mathbf{u})}{\partial u_1} & \frac{\partial f_2(\mathbf{u})}{\partial u_2} & \frac{\partial f_2(\mathbf{u})}{\partial u_3} \end{bmatrix} = \begin{bmatrix} 2 & 4 & 6 \\ 2u_1u_2^3u_3^4 & 3u_1^2u_2^2u_3^4 & 4u_1^2u_2^3u_3^3 \end{bmatrix}$$

$$67. \frac{\partial^2 f(\mathbf{u})}{\partial \mathbf{u} \partial \mathbf{u}'} = \begin{bmatrix} \frac{\partial^2 f(\mathbf{u})}{\partial u_1^2} & \frac{\partial^2 f(\mathbf{u})}{\partial u_2 \partial u_1} & \frac{\partial^2 f(\mathbf{u})}{\partial u_3 \partial u_1} \\ \frac{\partial^2 f(\mathbf{u})}{\partial u_1 \partial u_2} & \frac{\partial^2 f(\mathbf{u})}{\partial u_2^2} & \frac{\partial^2 f(\mathbf{u})}{\partial u_3 \partial u_2} \\ \frac{\partial^2 f(\mathbf{u})}{\partial u_1 \partial u_3} & \frac{\partial^2 f(\mathbf{u})}{\partial u_2 \partial u_3} & \frac{\partial^2 f(\mathbf{u})}{\partial u_3^2} \end{bmatrix} = \begin{bmatrix} 2 & 0 & 0 \\ 0 & 0 & 4u_3^3 \\ 0 & 4u_3^3 & 12u_2u_3^2 \end{bmatrix}$$

68. When answering this question, it helps if you remember quadratic forms.

$$\mathbf{v}'\mathbf{A}\mathbf{v} = \begin{bmatrix} v_1 & v_2 & v_3 \end{bmatrix} \begin{bmatrix} 2 & 0 & 0 \\ 0 & 0 & 4u_3^3 \\ 0 & 4u_3^3 & 12u_2u_3^2 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = 2v_1^2 + 8u_3^3v_2v_3 + 12u_2u_3^2v_3^2$$

69. Plug the value given for \mathbf{u} into the above answer, yielding $\mathbf{v}'\mathbf{A}\mathbf{v} = 2v_1^2$. This is positive or zero for all vectors of real numbers \mathbf{v} , $\mathbf{v}'\mathbf{A}\mathbf{v} \geq 0$. (However $\mathbf{v}'\mathbf{A}\mathbf{v} > 0$ is

not correct, for example $\mathbf{v} = \begin{bmatrix} 0 \\ 1 \\ 2 \end{bmatrix} \neq \mathbf{0}$ yields $\mathbf{v}'\mathbf{A}\mathbf{v} = 0$.)

$$70. \text{The first order condition is } \frac{\partial f(\mathbf{u})}{\partial \mathbf{u}} = \begin{bmatrix} \frac{\partial f(\mathbf{u})}{\partial u_1} \\ \frac{\partial f(\mathbf{u})}{\partial u_2} \\ \frac{\partial f(\mathbf{u})}{\partial u_3} \end{bmatrix} = \begin{bmatrix} 2u_1 \\ u_3^4 \\ 4u_2u_3^3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}. \text{ Thus, } 2u_1 = 0,$$

$u_3^4 = 0$, and $4u_2u_3^3 = 0$. Solving yields $u_1 = 0$, $u_3 = 0$, and $u_2 = \text{anything}$. So for

any k , $\tilde{\mathbf{u}} = \begin{bmatrix} 0 \\ k \\ 0 \end{bmatrix}$ is a candidate to yield a minimum of the function. Letting \mathbf{A} denote

the Hessian matrix obtained previously, and plugging in $\mathbf{u} = \tilde{\mathbf{u}}$, yields $\mathbf{v}'\mathbf{A}\mathbf{v} = 2v_1^2$ as in the previous question. This is ≥ 0 for all $\mathbf{v} \neq \mathbf{0}$. However it is not > 0 for all $\mathbf{v} \neq \mathbf{0}$, as demonstrated in the previous answer. Hence form B of the second order condition cannot guarantee that $\tilde{\mathbf{u}}$ yields a local minimum of the function. Were $\mathbf{v}'\mathbf{A}\mathbf{v}$ sometimes positive and sometimes negative, then you would know that $\tilde{\mathbf{u}}$ yields neither a maximum nor a minimum of the function, but this is not true either. So in this case, form B of the second order condition cannot tell you whether or not

$\tilde{\mathbf{u}} = \begin{bmatrix} 0 \\ k \\ 0 \end{bmatrix}$ yields a minimum. (You can be sure it is not a maximum.) In fact it is not a

minimum; for example $f\left(\begin{bmatrix} 0 & k & 0 \end{bmatrix}'\right) = 0$ but $f\left(\begin{bmatrix} 0 & -2 & 2 \end{bmatrix}'\right) = -8$.